

Harmaline-resistant mutant of *Methanothermobacter thermotrophicus* with a lesion in Na⁺/H⁺ antiporter

Monika Vidová¹, Janette Bobálová² and Peter Šmigán¹

¹ Institute of Animal Biochemistry and Genetics, Slovak Academy of Sciences, Ivanka pri Dunaji, Slovak Republic

² Institute of Analytical Chemistry of the ASCR, v.v.i., Brno, Czech Republic

Abstract. A spontaneous mutant of *Methanothermobacter thermotrophicus* resistant to the Na⁺/H⁺ antiporter inhibitor harmaline was isolated. The Na⁺/H⁺ exchange activity in the mutant cells was remarkably decreased in comparison with wild-type cells. Na⁺/H⁺ antiporter activity of wild-type cells grown in the high Na⁺ concentration (125 mmol/l) was significantly increased as compared to the cells grown under low Na⁺ concentration (6.25 mmol/l) conditions. In contrast, harmaline resistant mutant showed almost the same Na⁺/H⁺ antiporter activity under both these conditions. While harmaline profoundly inhibited methanogenesis in the wild-type, increased methanogenesis was observed both in the presence and absence of harmaline in the mutant strain. ATP synthesis driven by methanogenic electron transport was significantly enhanced in the mutant cells. The experimental data revealed the differential expression of A flavoprotein and molybdenum-containing formylmethanofuran dehydrogenase 1 subunit C in harmaline-resistant mutant. The overexpression of these proteins might contribute to harmaline resistance. Taken together the results indicate that harmaline resistance in this mutant has arisen as a consequence of mutation(s) in antiporter gene(s) or protein(s) linked to antiporter activity. Moreover this work provides the evidence that Na⁺/H⁺ exchanger deficiency in harmaline-resistant mutant can induce overexpression of several proteins participating in methanogenesis.

Key words: Methanogens — Harmaline resistance — Sodium/proton exchange

Introduction

Methanogens as members of the *Archaea*, belong to a group of microorganisms containing both H⁺- and Na⁺-dependent energetic cycle and the Na⁺/H⁺ antiporter(s). Despite numerous studies, the physiological and bioenergetic roles of these two bioenergetic cycles and of the Na⁺/H⁺ antiporter have not been satisfactorily explained so far. Kaesler and Schönheit (1989) have suggested that primarily generated electrochemical sodium gradient ($\Delta\bar{\mu}_{\text{Na}^+}$) can be converted in *Methanothermobacter thermotrophicus* by means of Na⁺/H⁺ antiporter activity to electrochemical proton gradient ($\Delta\bar{\mu}_{\text{H}^+}$). The resulting $\Delta\bar{\mu}_{\text{H}^+}$ can be utilized *via* the A₁A₀ ATP synthase for ATP production (Kaesler and Schönheit

1989; Schäfer et al. 1999). In methanogens containing cytochromes, the H₂:heterodisulfide oxidoreductase system and F₄₂₀ H₂:heterodisulfide oxidoreductase system are associated with membrane and coupled to proton translocation (Deppenmeier and Müller 2008). On the contrary it was suggested that in methanogens without cytochromes, the corresponding H₂:heterodisulfide reductase is only loosely associated with cytoplasmic membrane and is probably not coupled to proton translocation (Thauer et al. 2008). In such a case Na⁺/H⁺ antiporter might be an exclusive $\Delta\bar{\mu}_{\text{H}^+}$ generator. These ideas present an unusual new energetic situation in methanogens and Na⁺/H⁺ antiporter might acts in this complex energetic system as a basic regulatory element especially in cell without cytochromes, such as *M. thermoautotrophicus*. Inhibitory studies of Na⁺/H⁺ exchanger in methanoarchaea suggested that it is involved in pH and pNa homeostasis but its function in bioenergetics of methanoarchaea has not been resolved yet.

The understanding of the function of Na⁺/H⁺ exchanger and its cooperation with above mentioned bioenergetic sub-

Correspondence to: Peter Šmigán, Institute of Animal Biochemistry and Genetics, Slovak Academy of Sciences, Moyzesova 61, 900 28 Ivanka pri Dunaji, Slovak Republic
E-mail: peter.smigan@savba.sk

systems requires new experimental tools. We suggested that genetic approach to modulation or elimination of Na^+/H^+ antiport in cells of *M. thermautotrophicus* might be suitable for these purposes. Recently we have isolated a spontaneous mutant of *M. thermautotrophicus* resistant to the Na^+/H^+ antiporter inhibitor amiloride (Šurín et al. 2007). The study of the Na^+/H^+ antiporter in the amiloride-resistant mutant has shown that this mutant exhibits very specific phenotypic characteristics like reduced Na^+/H^+ antiport and elevated production of methane and ATP. The molecular study of amiloride-resistant mutant did not allow us to assign a Na^+/H^+ antiporter activity to a product of a particular gene from the *M. thermautotrophicus* genome or to identify protein(s) participating in this complex process. Therefore we decided to isolate mutants resistant to harmaline as another inhibitor of Na^+/H^+ antiporter in methanoarchaea (Schönheit and Beimborn 1985; Šmigáň et al. 1995). Such a mutant could have alterations in Na^+/H^+ antiporter itself and could thus provide a tool to extend our knowledge on the properties and the physiological role of Na^+/H^+ antiporter and its bioenergetic function in *M. thermautotrophicus*. Moreover, the molecular study of such mutant could provide a very promising experimental approach for identification of the protein/gene for the Na^+/H^+ antiporter and could lead to new ideas on the molecular mechanism of energy coupling in methanoarchaea. This paper reports on a mutant of *M. thermautotrophicus* resistant to harmaline exhibiting a lesion in Na^+/H^+ antiporter activity accompanied by an increase in methane and ATP formation. Moreover, this study revealed the differential expression of several proteins that may contribute to harmaline resistance in the mutant strain.

Materials and Methods

Growth conditions and isolation of harmaline-resistant mutant

Growth conditions of the *M. thermautotrophicus* strain ΔH were as described earlier (Šmigáň et al. 1984). Solid media were prepared by the addition of 1% (w/v) Gelrite to the medium. All manipulations were performed under strict anaerobic conditions. For the isolation of harmaline-resistant mutant of *M. thermautotrophicus*, growth medium (Šmigáň et al. 1984) containing K_2CO_3 instead of Na_2CO_3 was supplemented with 6.25 mmol/l NaCl. To select harmaline-resistant mutants, cultures were grown for 24 h in liquid medium in the presence of 800 $\mu\text{mol/l}$ harmaline. Cells were washed in growth medium and 1×10^8 cells were plated directly in an anaerobic box (Forma Scientific, model 1024) on solid medium (pH 7) containing harmaline. Plates were incubated at 60°C in a stainless-steel anaerobic jar for 14 days. Eight clones able to grow in the presence of harmaline were isolated and one of them was characterised in this study.

Measurements of Na^+/H^+ antiport activity

Na^+ -induced pH changes in weakly buffered cell suspensions were monitored by Orion 91-03 semimicroelectrode connected to Orion Research Ion Analyser 901 equipped with Kipp-Zonen BD40 recorder.

Spectrofluorimetric measurements of Na^+/H^+ antiport activity were performed as described earlier (Šmigáň et al. 1995) with following modifications. Exponentially grown cells were harvested and washed in 20 mmol/l MES buffer containing 50 mmol/l NH_4Cl (pH 6), under N_2 atmosphere, resuspended in 20 mmol/l HEPES, 10 mmol/l KCl (pH 7), to final concentration of 4.5 mg of cell proteins per milliliter, and maintained at 7°C under N_2 atmosphere. Aliquots of cells corresponding to 45 μg of cell protein were transferred to a stirred anaerobic cuvette containing 2 ml of buffer: 50 mmol/l HEPES, 10 mmol/l KCl, 5 mmol/l MgCl_2 and 2.5 mmol/l acridine orange preheated to 55°C. Quenching and Na^+ -induced dequenching of acridine orange were monitored under N_2 atmosphere at 55°C using Perkin-Elmer LS-50B spectrofluorimeter at excitation and emission wavelengths of 493 and 530 nm, respectively. Fluorescence before cells addition was assumed to be 100%.

Subcellular fractionation, analysis and identification of proteins

Soluble supernatant fractions (centrifuged at $100\,000 \times g$) and membrane vesicle proteins were isolated as described previously (Majerník et al. 2003). Membrane vesicles were also used to prepare chloroform/methanol extracts according to (Ruppert et al. 2001). Soluble supernatant fractions and membrane vesicles were analysed by a native polyacrylamide gel electrophoresis (PAGE) (Laemmli 1970) without the addition of sodium dodecyl sulfate (SDS) and 2-mercaptoethanol to buffers. SDS-PAGE was performed according to (Schägger and von Jagow 1987). Protein bands stained by Coomassie blue staining were excised from the gel, digested by trypsin and analysed using a MALDI TOF/TOF 4700 Proteomic Analyser (ABI). The Mascot search engine was used to identify cognate proteins by matching MS of detected peptides to *M. thermautotrophicus* proteins. Proteins were quantified by the Lowry method (Lowry et al. 1951) using bovine serum albumin (BSA) as standard.

Measurement of methane and ATP formation

Methane formation from CO_2 and H_2 by the cell suspension was measured by an analysis of glass vial headspace gases by GC. Aliquots were measured using a CarloErba Fractovap 4200 equipped with a 2-m steel column packed with Sepharon AE 200–300 μm and a thermal conductivity detector using argon as a carrier gas (Šmigáň et al. 1984). ATP synthesis




Figure 1. Changes of external pH in suspensions of *M. thermautotrophicus* wild-type (WT) cells (A) and harmaline resistant mutant (M) cells (B). Cells grown to exponential phase either in the presence of 125 mmol/l NaCl (high Na^+) or 6.5 mmol/l NaCl (low Na^+) were harvested, washed and resuspended in buffer (0.2 mol/l KCl, 2 mmol/l Tris : HCl, pH 6.8) to a concentration of 1.1 mg protein/ml, transferred to a double-necked vial and incubated at 60°C under constant stirring. **A.** High Na^+ cells: control (1), 800 $\mu\text{mol/l}$ harmaline (2). Low Na^+ cells: control (3), 800 $\mu\text{mol/l}$ harmaline (4). **B.** High Na^+ cells: control (1), 800 $\mu\text{mol/l}$ harmaline (2). Low Na^+ cells: control (3), 800 $\mu\text{mol/l}$ harmaline (4). Cells grown in the presence of 6.5 mmol/l NaCl. The arrows indicate the addition of NaCl to a final concentration of 60 mmol/l. All experiments were reproducibly repeated at least three times, and a representative data set is shown.

driven by methanogenic electron transport in cell suspensions of the wild-type and harmaline-resistant mutant was measured according to Schönheit and Perski (1983).

Results

Harmaline and amiloride have been shown to be competitive inhibitors of the Na^+/H^+ antiporter in *M. thermautotrophicus* (Schönheit and Beimborn 1985). It is important to emphasize that these two substances are chemically unrelated. Recently we have isolated a spontaneous mutant of *M. thermautotrophicus* resistant to the Na^+/H^+ antiporter inhibitor amiloride (Šurín et al. 2007).

To extend our knowledge on the function of this antiporter we tried to isolate a new spontaneous mutant of *M. thermautotrophicus* resistant to 800 $\mu\text{mol/l}$ harmaline.

Wild-type and the harmaline-resistant mutant strains showed comparable growth yields (0.245 mg and 0.24 mg of total cell protein/ml of growth medium, respectively) after 24 h of growth in standard culture media with 125 mmol/l NaCl. We hypothesized that harmaline-resistant mutant might have alterations in Na^+/H^+ antiporter itself as it was found in amiloride resistant mutant. To test the Na^+/H^+ antiporter activity of harmaline-resistant mutant and wild-type cells, the extent of Na^+ -induced acidification of the external medium by cell suspension cultivated in liquid media supplemented with 6.25 mmol/l or 125 mmol/l NaCl was monitored. The Na^+/H^+ exchanger activity in the mutant cells was remarkably decreased in comparison with wild-type cells. The Na^+/H^+ antiport activity of wild-type cells grown in the high sodium ion concentration (125 mmol/l) was significantly increased as compared to the cells grown under low Na^+ (6.25 mmol/l) conditions. On the other hand, the harmaline-resistant mutant showed almost the same Na^+/H^+ antiport activity under both conditions (Figure 1). Harmaline profoundly inhibited the Na^+/H^+ antiporter activity in the wild-type while this inhibitor even stimulated Na^+/H^+ exchange in mutant cells (Figure 1). The results of these experiments demonstrate that the increasing Na^+ ion concentration in growth medium leads to a significant increase of the Na^+/H^+ antiporter activity in wild-type cells. These results also indicate that Na^+/H^+ antiporter is inducible in *M. thermautotrophicus*.

To characterize Na^+/H^+ antiporter activity in the harmaline-resistant mutant strain in more detail, a spectrofluorimetric method was used for determination of kinetic parameters for the Na^+/H^+ exchange in both wild-type and mutant cells, respectively. The apparent maximum velocity V_{max} and Michaelis constant K_{M} were 73.6 a.u. (arbitrary unit) and 5.5 mmol/l for the wild-type, and 47.6 a.u. and 4.4 mmol/l for the mutant. Almost two-fold decrease in V_{max} in the mutant strain indicates either a reduced amount of Na^+/H^+ antiporter or a lower catalytic efficiency rather than a change in its affinity for Na^+ ions. These results were similar to our results obtained in the study of amiloride resistant mutant.

In 1985, Schönheit and Beimborn presented the evidence for a function of the Na^+/H^+ antiporter in methane formation and intracellular pH regulation (Schönheit and Beimborn 1985). It was therefore interesting to study the effect of significantly reduced activity of the Na^+/H^+ antiporter in harmaline-resistant mutant on Na^+ -dependent methane formation. As can be seen in Figure 2, harmaline-resistant mutant exhibited methanogenesis under resting conditions even in the presence of 800 $\mu\text{mol/l}$ harmaline, whereas methanogenesis was strongly inhibited under the same conditions in the wild-type cells. Since Na^+/H^+ antiporter activity is supposed to be sensitive to uncoupling, we tested the effect of the uncoupler 3,30,40,5-tetrachlorosalicylanilide




Figure 2. Time course of methane formation from H₂/CO₂ in the wild-type and harmfuline-resistant mutant of *M. thermautotrophicus*. Cells were suspended in 50 mmol/l MOPS-Tris buffer (pH 6.8), 10 mmol/l NaCl and 5 mmol/l MgCl₂ to a concentration of 1.8 mg protein/ml. The suspensions in 22 ml flasks were pressurized with H₂/CO₂ (4 : 1) to 100 kPa and incubated at 60°C. **A.** Wild-type cells: control (●), 800 μmol/l harmfuline (○) **B.** Mutant cells: control (▲), 800 μmol/l harmfuline (△). Methane was measured at the times indicated. All experiments were reproducibly repeated at least three times and a representative data set is shown.

(TCS) on growth and methane production in the wild-type and harmfuline-resistant mutant cells. Significant differences were found between the ability of mutant and wild-type cells to synthesize methane and to grow in the presence of this uncoupler (Table 1).

While harmfuline and TCS strongly inhibited growth and methanogenesis in the wild-type, both parameters were insensitive to these inhibitors in the mutant strain. These findings support the idea that in wild-type cells, $\Delta\bar{\mu}_{\text{Na}^+}$ can be converted to $\Delta\bar{\mu}_{\text{H}^+}$ by Na⁺/H⁺ antiporter (Kaesler and Schönheit 1989) and resulting $\Delta\bar{\mu}_{\text{H}^+}$ can be then directly utilized *via* the A₁A₀ ATP synthase for ATP production. In the mutant strain with modified Na⁺/H⁺ antiporter system, $\Delta\bar{\mu}_{\text{Na}^+}$ cannot be converted by means of the Na⁺/H⁺ antiporter to $\Delta\bar{\mu}_{\text{H}^+}$. Moreover, $\Delta\bar{\mu}_{\text{H}^+}$ cannot be generated in these cells by $\Delta\bar{\mu}_{\text{H}^+}$ generator – H₂: heterodisulphid reductase since this enzyme does not work in the methanoarchaea cells lacking cytochromes (Deppenmeier et al. 1996; Thauer et al. 2008). This indicates that Na⁺ ions are the exclusive coupling ions in this mutant. Amiloride (800 μmol/l), another inhibitor of the Na⁺/H⁺ antiporter in methanoarchaea, inhibited methane formation in the harmfuline-resistant mutant strain under non-growing conditions (not shown). This demonstrated the absence of crossresistance between harmfuline and amiloride in the harmfuline-resistant mutant and indicated different biochemical mechanisms of re-

sistance to harmfuline and amiloride in *M. thermautotrophicus*. Na⁺/H⁺ antiporter in methanoarchaea seems to be a process that couples the circulation of H⁺ and Na⁺ ions across the cytoplasmic membrane.

As mentioned earlier, Na⁺/H⁺ antiporter could modulate the driving force for ATP synthesis under physiological conditions (Kaesler and Schönheit 1989; Šmigáň et al. 1995). Therefore, we studied the potential role of the Na⁺/H⁺ antiporter in total ATP synthesis in more detail. Comparison of wild-type and mutant ATP synthesis driven by methanogenic electron transport has shown that total ATP synthesis is

Table 1. Effect of TCS and harmfuline on methane production of wild-type and harmfuline-resistant mutant^a

Treatment	CH ₄ /24 h (mmol)	
	Wild-type	Mutant
Control	1.3	1.2
TCS	0.26	1.2
Harmaline	0.0	1.16

^a Wild-type and harmfuline-resistant mutant were cultivated in medium supplemented with 125 mmol/l NaCl in the presence of 3 μmol/l TCS or 800 μmol/l harmfuline. Methane was determined after 24 h by gas chromatography.

higher in the mutant strain (Figure 3). We have also determined the effect of significantly reduced Na^+/H^+ antiporter activity in the mutant strain on ATP synthesis driven by an electrogenic potassium efflux in the presence of sodium ions. The results clearly showed significantly enhanced ATP synthesis in the harmaline-resistant mutant (Figure 4). It should be noted in this context that native PAGE combined with MALDI TOF/TOF mass spectrometry did not reveal increased levels of A_1 and A_0 subcomplexes of the A_1A_0 -type synthase in the mutant strain (not shown).

These findings indicate that the Na^+/H^+ antiporter may compete with the Na^+ -translocating ATP synthase for sodium ions during ATP synthesis, as it was suggested previously (Šmigáň et al. 1995). Similarly, Becher and Müller (1994) have found that inhibition of the Na^+/H^+ antiporter in *M. mazei* led to a stimulation of ATP synthesis driven by a methyl transfer reaction generating $\Delta\mu_{\text{Na}^+}$, as well as by artificially created $\Delta p\text{Na}$.

To identify molecular basis of the harmaline resistance, mass spectrometry analysis of the protein composition of membrane vesicles and supernatant fraction derived from wild-type and

harmaline-resistant cells was performed. The preliminary results of MS analysis of the protein composition of membrane vesicles and supernatant fractions revealed that mutant cells contained significantly higher amounts of type A flavoprotein and molybdenum-containing formylmethanofuran dehydrogenase. It has been suggested recently that these proteins play a direct role in flavin-based electron bifurcation in methanoarchaea lacking cytochromes (Thauer et al. 2008).

Discussion

The coexistence of two primary ion gradients $\Delta\tilde{\mu}_{\text{H}^+}$ and $\Delta\tilde{\mu}_{\text{Na}^+}$ in methanoarchaea representing a completely unprecedented energetic situation in prokaryotic cells could have lead to the evolution of specific transport system(s). This system might be a basic regulatory element responsible for the homeostasis of Na^+ - and H^+ -cycles in methanoarchaea. We suggested Na^+/H^+ antiporter as one of candidates for this role. To test this hypothesis, we applied genetic approach to prepare a model system in which modulation and/or elimination of this antiporter would be viable.

Three years ago we isolated a spontaneous *M. thermotrophicus* mutant resistant to the Na^+/H^+ antiporter inhibitor amiloride (Šurín et al. 2007). Biochemical and proteomic study of this mutant did not definitely clarify neither the predicted function of this antiporter in balancing the energy of $\Delta\tilde{\mu}_{\text{H}^+}$ and $\Delta\tilde{\mu}_{\text{Na}^+}$ nor the protein components responsible for Na^+/H^+ exchange in the bioenergetics of *M. thermotrophicus*. As an extension of these studies we isolated harmaline-resistant mutant of *M. thermotrophicus*. The finding that growth in the presence of high sodium concentration is associated with an elevated Na^+/H^+ antiporter activity in wild-type cells supports the idea that the expression of this exchanger is regulated by sodium ions. It also indicates that sodium dependence of methanoarchaea is mediated not only by N-5-methyltetrahydromethanopterin: coenzyme M methyltransferase (EC 2.1.1.86) as generator of $\Delta\tilde{\mu}_{\text{Na}^+}$ (a primary sodium pump) (Becher et al. 1992) but also by the Na^+/H^+ exchanger. Recently we observed a significant increase in Na^+/H^+ antiporter activity when cells progressed into the stationary growth phase. We can assume that *M. thermotrophicus* has either one Na^+/H^+ antiporter differently regulated by Na^+ concentration or that there are at least two independent Na^+/H^+ antiporter systems each controlled by different mechanisms. Results obtained from kinetic measurements indicate that Na^+/H^+ antiporter in the mutant strain is either present in reduced amounts or it exhibits lower catalytic efficiency rather than it would have modified affinity for Na^+ ions. Similar results were achieved also in the study of amiloride resistant mutant. As mentioned above, *M. thermotrophicus* belongs to a group of methanoarchaea without cytochromes, and its




Figure 3. Methanogenesis-driven ATP synthesis in the wild-type and harmaline-resistant mutant of *M. thermotrophicus*. Cells were suspended in 50 mmol/l MOPS-Tris buffer (pH 6.8), 10 mmol/l NaCl and 5 mmol/l MgCl_2 to a concentration of 1.1 mg protein/ml. Gas phase was 80% H_2 /20% CO_2 . ● wild-type cells, ○ mutant cells. All experiments were reproducibly repeated at least three times and a representative data set is shown.

H₂:heterodisulphid reductase is a cytoplasmic enzyme and is not coupled to transmembrane proton translocation. If this assumption is valid, Na⁺/H⁺ antiporter is the only possible generator of $\Delta\bar{\mu}_{H^+}$.

The results described above have raised the question if harmaline-resistant mutant cells can grow when $\Delta\bar{\mu}_{H^+}$ is dissipated in the presence of protonophore TCS. To answer this question, we studied the inhibitory effect of TCS on the methane formation in growing wild-type and harmaline-resistant mutant cells (Table 1). Our results indicate that $\Delta\bar{\mu}_{H^+}$ is a driving force for at least some endergonic processes in the wild-type cells grown in the presence of high NaCl (125.0 mmol/l NaCl) and that Na⁺/H⁺ exchange activity sensitive to harmaline is responsible for generation of $\Delta\bar{\mu}_{H^+}$. On the other hand, mutant cells can grow under the same conditions in the presence of protonophore TCS and harmaline. This supports the idea that harmaline-resistant mutant with modified Na⁺/H⁺ antiporter can grow in the absence of $\Delta\bar{\mu}_{H^+}$ that should be dissipated in the presence of uncoupler. This finding also indicates that only




Figure 4. Potassium diffusion potential-driven ATP synthesis in the wild-type and harmaline-resistant mutant of *M. thermautotrophicus*. Cells were suspended in 50 mmol/l MOPS-Tris buffer (pH 6.8) to a concentration of 3.0 mg protein/ml. After 5 min of preincubation at 60°C, ATP synthesis was induced by the addition of valinomycin (f.c. 15 μmol/l). At $t = 5$ min, NaCl was added to f.c. 50 mmol/l. Wild-type cells: control (●), 3 μmol/l TCS (○). Mutant cells: control (▲), 3 μmol/l TCS (△).

$\Delta\bar{\mu}_{Na^+}$ generated by N-5-methyltetrahydromethanopterin: coenzyme M methyltransferase (EC 2.1.1.86) drives ATP synthesis and that Na⁺ ion is the exclusive coupling ion in this mutant. This means also that $\Delta\bar{\mu}_{H^+}$ generator H₂: heterodisulphid reductase, generating $\Delta\bar{\mu}_{H^+}$ by redox potential-driven proton translocation (Deppenmeier et al. 1996), does not work in cells lacking cytochromes as suggested by Thauer et al. (2008). These authors suggested recently that in methanoarchaea group with cytochromes the first and last steps in methanogenesis from H₂ and CO₂ are coupled chemiosmotically (Thauer et al. 2008). *M. thermautotrophicus* belong to a group lacking cytochromes. Gunsalus and Wolfe (1977) observed in *M. thermautotrophicus* a stimulation of CO₂ reduction to methane by methyl-coenzyme M. It was suggested that the first and last steps of methanogenesis in these methanoarchaea are energetically coupled by a cytoplasmic enzyme complex that mediates flavin-based electron bifurcation (Thauer et al. 2008). The study of harmaline-resistant mutant opened the possibility to study the role of Na⁺/H⁺ antiporter in these complex bioenergetic systems. Therefore we attempted to define on the protein level the molecular basis of the harmaline resistance and of accompanying phenotypic characteristics of the mutant strain. We compared the level of protein expression in different cellular fractions. The preliminary results of mass spectrometry analysis of membrane proteins in wild-type and harmaline-resistant cells revealed that the mutant strain contained significantly higher amounts of type A flavoprotein and molybdenum-containing formylmethanofuran dehydrogenase 1 subunit C. Low concentrations or absence of the Na⁺/H⁺ antiporter in harmaline-resistant mutant might be compensated for by elevated amounts of type A flavoprotein and molybdenum-containing formylmethanofuran dehydrogenase participating in flavin-based electron bifurcation. All these findings support the idea that Na⁺/H⁺ exchanger might be a clue to the adaptive bioenergetic behaviour of *M. thermautotrophicus* under the different physiological conditions. Moreover this work provides the evidence that Na⁺/H⁺ exchanger deficiency in harmaline-resistant mutant can induce changes at the expression level of several genes participating in methanogenesis. The results imply that harmaline resistance is a consequence of decreased Na⁺/H⁺ antiporter expression.

Acknowledgements. The authors gratefully acknowledge the technical assistance of Mrs. Petra Melicherová and Mrs. Lenka Országhová. This investigation was supported in part by the Slovak Research and Development Agency APVT-51-024904, APVV-VVCE 0064-07 and by research grants VEGA 2/0015/09 from the Slovak Academy of Sciences and the Institutional Research Plan AV0Z40310501 of the Institute of Analytical Chemistry of the ASCR, v.v.i. M. V., J. B. and P. Š. have no conflict of interest and no financial interest in the publication of this manuscript.

References

- Becher B., Müller V. (1994): $\Delta\mu_{\text{Na}^+}$ drives the synthesis of ATP via an $\Delta\mu_{\text{Na}^+}$ -translocating F_1F_0 -ATP synthase in membrane vesicles of the archaeon *Methanosarcina mazei* Gö1. *J. Bacteriol.* **176**, 2543–2550
- Becher B., Müller V., Gottschalk G. (1992): The methyl-tetrahydromethanopterin:coenzyme M-methyltransferase of *Methanosarcina* strain Gö1 is a primary sodium pump. *FEMS Microbiol. Lett.* **91**, 239–244
- Deppenmeier U., Müller V., Gottschalk G. (1996): Pathways of energy conservation in methanogenic archaea. *Arch. Microbiol.* **165**, 149–163
doi:10.1007/BF01692856
- Deppenmeier U., Müller V. (2008): Life close to the thermodynamic limit: how methanogenic archaea conserve energy. *Results and Probl. Cell Differ.* **45**, 123–152
doi:10.1007/400_2006_026
- Gunsalus R. P., Wolfe R. S. (1977): Stimulation of CO₂ reduction to methane by methyl-coenzyme M in extracts of *Methanobacterium*. *Biochem. Biophys. Res. Commun.* **76**, 790–795
doi:10.1016/0006-291X(77)91570-4
- Kaesler B., Schönheit P. (1989): The sodium cycle in methanogenesis. CO₂ reduction to the formaldehyde level in methanogenic bacteria is driven by a primary electrochemical potential of Na⁺ generated by formaldehyde reduction to CH₄. *Eur. J. Biochem.* **186**, 309–316
doi:10.1111/j.1432-1033.1989.tb15210.x
- Laemmli U. K. (1970): Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* **227**, 680–685
doi:10.1038/227680a0
- Lowry O. H., Rosebrough N. J., Farr A. L., Randall R. J. (1951): Protein measurement with the Folin phenol reagent. *J. Biol. Chem.* **193**, 265–275
- Majerník A., Čuboňová L., Polák P., Šmigáň P., Greksák M. (2003): Biochemical analysis of neomycin-resistance in the methanoarchaeon *Methanothermobacter thermoautotrophicus* and some implications for energetic processes in this strain. *Anaerobe* **9**, 31–38
doi:10.1016/S1075-9964(03)00042-8
- Ruppert C., Schmid R., Hedderich R., Müller V. (2001): Selective extraction of subunit D of the Na⁺-translocating methyltransferase and subunit c of the A₁A₀ ATPase from the cytoplasmic membrane of methanogenic archaea by chloroform/methanol and characterization of subunit c of *Methanothermobacter thermoautotrophicus* as a 16-kDa proteolipid. *FEMS Microbiol. Lett.* **195**, 47–51
doi:10.1111/j.1574-6968.2001.tb10496.x
- Schägger H., von Jagow G. (1987): Tricine-sodium dodecyl sulfate-polyacrylamide gel electrophoresis for the separation of proteins in the range from 1 to 100 kDa. *Anal. Biochem.* **166**, 368–379
doi:10.1016/0003-2697(87)90587-2
- Schäfer G., Engelhard M., Müller V. (1999). Bioenergetics of the Archaea. *Microbiol. Mol. Biol. Rev.* **63**, 570–620
- Schönheit P., Beimborn D. B. (1985): Presence of Na⁺/H⁺ antiporter in *Methanobacterium thermoautotrophicum* and its role in Na⁺ dependent methanogenesis. *Arch. Microbiol.* **142**, 354–361
doi:10.1007/BF00491903
- Schönheit P., Perski H. J. (1983): ATP synthesis driven by potassium diffusion potential in *Methanobacterium thermoautotrophicum* is stimulated by sodium. *FEMS Microbiol. Lett.* **20**, 263–267
doi:10.1111/j.1574-6968.1983.tb00129.x
- Šmigáň P., Friederová A., Rusnák P., Greksák M. (1984): Effect of 2,4-dinitrophenol and ionophores on growth and methanogenesis in *Methanobacterium thermoautotrophicum*. *Folia Microbiol.* **29**, 353–358
doi:10.1007/BF02887762
- Šmigáň P., Majerník A., Polák P., Hapala I., Greksák M. (1995): The presence of H⁺ and Na⁺-translocating ATPases in *Methanobacterium thermoautotrophicum* and their possible function under alkaline conditions. *FEBS Lett.* **371**, 119–122
doi:10.1016/0014-5793(95)00866-8
- Šurín S., Čuboňová L., Majerník A., McDermott P., Chong J. P. J., Šmigáň P. (2007): Isolation and characterization of an amiloride-resistant mutant of *Methanothermobacter thermoautotrophicus* possessing a defective Na⁺/H⁺ antiport. *FEMS Microbiol. Lett.* **269**, 301–308
doi:10.1111/j.1574-6968.2007.00655.x
- Thauer K. R., Kaster A. K., Seedorf H., Buckel W., Hedderich R. (2008): Methanogenic archaea: ecologically relevant differences in energy conservation. *Nature Rev. Microbiol.* **6**, 579–591
doi:10.1038/nrmicro1931

Received: October 19, 2010

Final version accepted: November 28, 2010